MAT 223, Discussion Questions 11.16

1. What conditions need to be satisfied to use the normal approximation for proportions?

and n(1-p)>5

2. Find a 95% confidence interval for $\hat{p}=59\%$ and a sample size of 1100.

1 Prop 2 Int

(.56094, 61906)

X=.59*1100 = 649 6 must be a whole #

- 1) = 1100 C-level: .95

 3. How big a sample size is needed to estimate a true proportion within 1%? (The most conservative estimate for p, if it is not known, is to assume a value of 50%.)

$$n = \left(\frac{2\omega h}{E}\right)^2 p(1-p) = \left(\frac{1.9b}{.01}\right)^2 (.5) (5) = 9604$$

4. The student newspaper at a college asks a simple random sample of 250 undergraduates, "Do you favor eliminating supplemental fees for lab courses?" In all, 150 of the 250 are in favor of eliminating such fees. Find a 95% confidence interval not using the quick method.

(Prop ZIht

(.53927, .66073)

X = 150 n=250

C-level: 95

5. A recent Gallup Poll interviewed a random sample of 1523 adults. Of these, 868 bought a lottery ticket in the past year. Construct a 99% confidence interval for these results.

1 Prop Z. Int

N=1523

(.41759, .48327)

- Clovel: 99
- 6. A poll of 1234 adults found that 62% expect an increase in environmental pollution in the next decade. Take the poll's sample to be a simple random sample of all adults. What is an 98% confidence interval for these results?

1 Prop & Int

X=, 62*1234 = 765.08 => 765 (must be a whole #)

N= 1234

C-level: 98

(.58779, .65208)

7. Read and comment on the attached article about learning strategies.			
THE POST WWW. HE WALLETTING COMPARTICIO (AND 1/107 Lynn) II - II 40			
learning/?full=true&prir	nt=true#.VRbbVI7F-p0		ets or successiul-