

Instructions: Work the problems below as directed. Show all work. Clearly mark your final answers. Use exact values unless the problem specifically directs you to round. Simplify as much as possible. Partial credit is possible, but solutions without work will not receive full credit.

1. Find the indicated derivative of the given function.

a. $f'(x), f(x) = \sqrt{b + \sec \pi x^2} = (b + \sec \pi x^2)^{1/2}$

$$f'(x) = \frac{1}{2}(b + \sec \pi x^2)^{-1/2} (2x \sec \pi x^2 \tan \pi x^2 \cdot \pi) = \frac{\pi x \sec \pi x^2 \tan \pi x^2}{\sqrt{b + \sec \pi x^2}}$$

b. $f'(x), f(x) = \tan(\sec x)$

$$f'(x) = \sec^2(\sec x) \cdot \sec x \tan x$$

c. $f''(x), f(x) = (3x - 2)^6$

$$f'(x) = 6(3x - 2)^5 (3) = 18(3x - 2)^5$$

$$f''(x) = 18 \cdot 5(3x - 2)^4 (3) = 270(3x - 2)^4$$

d. $f''(x), f(x) = x \sin x$

$$f'(x) = \sin x + x \cos x$$

$$f''(x) = \cos x + \cos x - x \sin x = 2\cos x - x \sin x$$

2. A particle is moving along a curve with a velocity of $v(t) = \frac{1}{3}t^3 - 2t^2 + 6t$. Find the acceleration of the particle at any time t . When is the acceleration of the particle zero?

$$v'(t) = a(t) = t^2 - 4t + 6$$

$$t^2 - 4t + 6 = 0$$

$$t = \frac{4 \pm \sqrt{16 - 4(1)(6)}}{2(1)} = \frac{4 \pm \sqrt{16 - 24}}{2}$$

The acceleration is never zero.